

# E-S-C-M

## Evaluation of the Socially interactive Communication in Mutism

First name: \_\_\_\_\_ 

m	f
---	---

 Date: \_\_\_\_\_

Name: \_\_\_\_\_ First evaluation:

Date of birth: \_\_\_\_\_ Age: \_\_\_\_\_ Next evaluation:  Nr.: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ Investigator: \_\_\_\_\_

0 = not stressful: uninhibited communication  
 1 = moderately stressful: communication on request  
 2 = very stressful: selective mutism

Socially interactive situation	Communicative load factor	0	1	2
Use of facial expressions and gestures		0	0	0
Talking at home with parents and siblings		0	0	0
Talking to grandparents and other relatives		0	0	0
Answering questions (adults) from neighbors		0	0	0
Playing and talking with neighbor children		0	0	0
Playing and talking with kids in kindergarten or at school		0	0	0
Communicative contact to educators and teachers		0	0	0
Communicative contact to work colleagues		0	0	0
Communication via literary language		0	0	0
Sounds with the mouth at home		0	0	0
Sounds with the mouth in the therapeutic setting		0	0	0
A-word-answers outside the family		0	0	0
A-phrase-answers outside the family		0	0	0
Singing at home		0	0	0
Singing in the therapeutic setting		0	0	0
Greetings and goodbyes		0	0	0
Retelling stories at home		0	0	0
Retelling stories outside the family		0	0	0
Making phone calls		0	0	0
Speaking in therapeutic accompaniment outside the practice/in-vivo-therapy		0	0	0
Free conversation in the therapy situation		0	0	0
Independent shopping with conversation		0	0	0
Question-answer patterns with strangers		0	0	0

∑ Next evaluation

∑ First evaluation